

CIRIC TYPE COINCIDENCE & FIXED POINT RESULTS FOR NON EXPANSIVE SINGLE VALUED MAPS IN 2-METRIC SPACES

Sukh Raj Singh¹, R. D. Daheriya², Manoj Ughade³

¹Department of Mathematics (Research Scholar),

J.H. Govt. Post Graduate College, Betul, India-460001

²Department of Mathematics, J.H. Govt. Post Graduate College, Betul, India-460001

³Department of Mathematics, Sarvepalli Radhakrishnan University, Bhopal, India-462026

ABSTRACT

In this paper, we consider the existence of coincidences and fixed points of non-expansive type conditions satisfied by single valued maps and prove some fixed point theorems for non-expansive type single valued mappings in 2-metric space.

2000 HEMATICS SUBJECT CLASSIFICATIONS: 47H10, 54H25

Keywords: 2-Metric Spaces; Coincidence Point; Common Fixed Point; Compatible Mappings; Non-Expansive Mappings.

I INTRODUCTION

Fixed point theorems for contractive, non-expansive, contractive type and non-expansive type mappings provide techniques for solving a variety of applied problems in mathematical and engineering sciences. It is one of the reason that many authors have studied various classes of contractive type or non-expansive type mappings. If T is such that for all x, y in X

$$(1.1) \quad d(Tx, Ty) \leq \lambda d(x, y)$$

where $0 < \lambda < 1$, then T is said to be a contraction mapping. If T satisfies (1.1) with $\lambda = 1$, then T is called a non-expansive mapping. If T satisfies any conditions of type

$$(1.2) \quad d(Tx, Ty) \leq a_1 d(x, y) + a_2 d(x, Tx) + a_3 d(y, Ty) + a_4 d(x, Ty) + a_5 d(y, Tx)$$

where a_i ($i = 1, 2, 3, 4, 5$) are nonnegative real numbers such that $a_1 + a_2 + a_3 + a_4 + a_5 < 1$, then T is said to be a contractive type mapping. If T satisfies (1.2) with $a_1 + a_2 + a_3 + a_4 + a_5 = 1$, then T is said to be a non-expansive type mapping. Similar terminology is used for multi-valued mappings.

Bogin [3] proved the following result:

Theorem 1.1 Let X be a nonempty complete metric space and $T: X \rightarrow X$ a mapping satisfying

$$(1.3) \quad d(Tx, Ty) \leq ad(x, y) + b[d(x, Tx) + d(y, Ty)] + c[d(x, Ty) + d(y, Tx)]$$

where $a \geq 0, b > 0, c > 0$ and

$$(1.4) \quad a + 2b + 2c = 1$$

Then T has a unique fixed point.

This result was generalized by Rhoades [19] and Ćirić [6,7]. Iseki [13] studied a family of commuting mappings T_1, T_2, \dots, T_n which satisfy (1.3) with $a \geq 0, b \geq 0, c \geq 0$ and $a + 2b + 2c = 1$. For Banach spaces the famous is Gregus's Fixed Point Theorem [11] for non-expansive type single-valued mappings, which satisfy (1.3) with $c = 0, a < 1$. Ćirić [6] introduced and investigated a new class of self-mappings T on X which satisfy an inequality of type (1.3) with $b \geq 0$ and still have a fixed point. Also proved that by an example if the mapping T satisfies (1.3) with $b = 0$ and if a and c are such that (1.4) holds, then T need not have a fixed point. Therefore, a contractive condition for T , which shall guarantee a fixed point of T in the case $b = 0$ and $a + 2c = 1$, must be stricter than (1.3).

The concept of 2-metric space is a natural generalization of the classical one of metric space. It has been investigated, initially, by Gähler and has been developed extensively by Gähler and many other mathematicians [8-10]. The topology induced by 2-metric space is called 2-metric topology, which is generated by the set of all open spheres with two centers. Iseki [12] studied the fixed point theorems in 2-metric spaces. A number of fixed point theorems has been proved for 2-metric spaces. Liu and Zhang [15] proved a few necessary and sufficient conditions for the existence of a common fixed point of a pair of mappings in 2-metric spaces. These results have generalized and improved by a number of mathematicians. Singh, Adiga and Giniswami [20] proved a fixed point theorem in 2-metric spaces for non-expansive type mappings.

We recall the following definitions and results which can be found in [8].

Definition 2.1 (see [8]) Let X be a nonempty set. A real valued function d on X^3 is said to a 2-metric if, for all $x, y, z, u \in X$, the following conditions hold:

- (1) To each pair of distinct points x, y in X , there exists a point $z \in X$ such that $d(x, y, z) \neq 0$;

- (2) $d(x, y, z) = 0$ if at least two of x, y, z are equal;
- (3) $d(x, y, z) = d(x, z, y) = d(y, x, z) = d(y, z, x) = d(z, x, y) = d(z, y, x)$;
- (4) $d(x, y, z) \leq d(x, y, u) + d(x, u, z) + d(u, y, z)$.

Then (X, d) is called a 2-metric space which will be sometimes denoted by X if there is no confusion. Every member $x \in X$ is called a point in X . Geometrically a 2-metric $d(x, y, z)$ represents the area of a triangle with vertices x, y and z .

Definition 2.2 A sequence $\{x_n\}$ in a 2-metric space (X, d) is said to be convergent to a point $x \in X$, if $\lim_{n \rightarrow \infty} d(x_n, x, u) = 0$ for all $u \in X$.

Definition 2.3 A sequence $\{x_n\}$ in a 2-metric space (X, d) is said to be Cauchy sequence if for all $z \in X$, $\lim_{n \rightarrow \infty} d(x_n, x_m, z) = 0$.

Definition 2.4 A sequence $\{x_n\}$ in a 2-metric space (X, d) is said to be complete if every Cauchy sequence in X is convergent.

Definition 2.5 (see [16]) Let f and T be mappings from a 2-metric space (X, d) into itself. The pair (f, T) is said to be compatible pair (co. p.) if for all $u \in X$, $\lim_{n \rightarrow \infty} d(fTx_n, Tfx_n, u) = 0$ whenever $\{x_n\}$ is a sequence in X such that $\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} Tx_n = p$ for some $p \in X$.

Definition 2.6 (see [16]) Let f and T be mappings from a 2-metric space (X, d) into itself. The pair (f, T) is said to be weakly compatible pair (w. co. p.) if $fx = Tx$ (for some $x \in X$) implies $fTx = Tfx$.

Definition 2.7 (see [16]) Let f and T be mappings from a 2-metric space (X, d) into itself. The pair (f, T) is said to be compatible of type (A) if

$$\lim_{n \rightarrow \infty} d(Tfx_n, ffx_n, u) = \lim_{n \rightarrow \infty} d(fTx_n, TTx_n, u) = 0$$

for all $u \in X$, whenever $\{x_n\}$ is a sequence in X such that $\lim_{n \rightarrow \infty} Sx_n = \lim_{n \rightarrow \infty} Tx_n = p$ for some $p \in X$.

Definition 2.8 (see [16]) Let f and T be mappings from a 2-metric space (X, d) into itself. The pair (f, T) is said to be weakly compatible of type (A) if

$$\lim_{n \rightarrow \infty} d(fTx_n, TTx_n, z) \leq \lim_{n \rightarrow \infty} d(Tfx_n, TTx_n, z)$$

and

$$\lim_{n \rightarrow \infty} d(Tfx_n, ffx_n, z) \leq \lim_{n \rightarrow \infty} d(fTx_n, ffx_n, z)$$

for all $z \in X$, whenever $\{x_n\}$ is a sequence in X such that $\lim_{n \rightarrow \infty} Sx_n = \lim_{n \rightarrow \infty} Tx_n = t$ for some $t \in X$.

Example: 2.9 (see [17]) Define d on $[0,1] \times [0,1] \times [0,1]$ by $d(x, y, z) = \min\{\rho(x, y), \rho(y, z), \rho(z, x)\}$, where ρ is a usual metric on $[0,1]$. Then it is easy to see that d is a 2-metric on $[0,1]$. Define $f, T: [0,1] \rightarrow [0,1]$ by $fx = \frac{x}{1+x}$ and $Tx = \frac{x}{2}$. Choose a sequence $\{x_n\}$ in $[0,1]$ such that converges to zero in $[0,1]$ i.e. $x_n \rightarrow 0 \in [0,1]$ as $n \rightarrow \infty$. Then for all $u \in X$,

$$\begin{aligned} \lim_{n \rightarrow \infty} d(fTx_n, Tfx_n, u) &= \lim_{n \rightarrow \infty} d\left(f\left(\frac{x_n}{2}\right), T\left(\frac{x_n}{1+x_n}\right), u\right) \\ &= \lim_{n \rightarrow \infty} d\left(\frac{\left(\frac{x_n}{2}\right)}{1+\left(\frac{x_n}{2}\right)}, \frac{\left(\frac{x_n}{1+x_n}\right)}{2}, u\right) \\ &= \lim_{n \rightarrow \infty} d\left(\frac{x_n}{2+x_n}, \frac{x_n}{2(1+x_n)}, u\right) \\ &= \lim_{n \rightarrow \infty} \min\left\{\rho\left(\frac{x_n}{2+x_n}, \frac{x_n}{2(1+x_n)}\right), \rho\left(\frac{x_n}{2(1+x_n)}, u\right), \rho\left(u, \frac{x_n}{2+x_n}\right)\right\} \\ &\leq \lim_{n \rightarrow \infty} \rho\left(\frac{x_n}{2+x_n}, \frac{x_n}{2(1+x_n)}\right) \\ &= \lim_{n \rightarrow \infty} x_n \left| \frac{1}{2+x_n} - \frac{1}{2(1+x_n)} \right| = 0. \end{aligned}$$

Thus, $d(fTx_n, Tfx_n, u) \rightarrow 0$, as $n \rightarrow +\infty$ when $x_n \rightarrow 0 \in X$ as $n \rightarrow +\infty$. Hence (f, T) is a co. p. In view of Proposition 2.4 of [18], every pair of compatible mappings of type (A) is weakly compatible mappings of type (A) whereas in view of Proposition 2.9 of [18], every pair of compatible mappings of type (A) is weakly compatible pair.

In this paper, we prove Ciric [6] type common fixed point theorems under non-expansive type conditions in the setting of 2-metric spaces. We shall investigate a class of self-mappings T, f on X which satisfy the following non-expansive type condition:

$$\begin{aligned} (1.5) \quad d(Tx, Ty, u) &\leq a(x, y) \max\{d(fx, fy, u), d(fx, Tx, u), d(fy, Ty, u), \frac{1}{2}[M(x, y, u) + m(x, y, u)]\} \\ &\quad + c(x, y)[M(x, y, u) + hm(x, y, u)] \end{aligned}$$

for all $x, y, u \in X$, where

$$M(x, y, u) = \max\{d(fx, Ty, u), d(fy, Tx, u)\}$$

$$m(x, y, u) = \min\{d(fx, Ty, u), d(fy, Tx, u)\}$$

and

$$(1.6) \quad 0 < h < 1, a(x, y) > 0, \beta = \inf \{c(x, y) : x, y \in X\} > 0$$

$$(1.7) \quad \sup_{x, y \in X} (a(x, y) + 2c(x, y)) = 1.$$

II MAIN RESULT

Now, we give our main results.

Theorem 2.1 Let (X, d) be a 2-metric space, T, f are self maps of X satisfying condition (1.5), where a and c satisfying (1.6) and (1.7) with $T(X) \subseteq f(X)$ and either (a) X is complete and f is surjective; or (b) X is complete, f is continuous and T, f are compatible; or (c) $f(X)$ is complete; or (d) $T(X)$ is complete. Then f and T have a coincidence point in X . Further, the coincidence value is unique, i.e. $fp = fq$ whenever $fp = Tp$ and $fq = Tq, (p, q \in X)$.

Proof Let $x_0 \in X$. We construct two sequences $\{x_n\}$ and $\{y_n\}$ as follows: Since $T(X) \subseteq f(X)$, choose x_1 so that $y_1 = fx_1 = Tx_0$. In general, choose x_{n+1} so that $y_{n+1} = fx_{n+1} = Tx_n, n \in \mathbb{N}$. For simplicity, we set $d_n(u) = d(y_n, y_{n+1}, u)$ for all $u \in X$ and $n \in \mathbb{N} \cup \{0\}$. Obviously, $d_n(y_n) = 0 = d_n(y_{n+1}) \forall n \in \mathbb{N} \cup \{0\}$.

First, we claim that $d_n(y_{n+2}) = 0$. On the contrary, suppose that $d_n(y_{n+2}) \neq 0, \forall n \in \mathbb{N} \cup \{0\}$. Applying (1.5), we have

$$\begin{aligned} (2.1) \quad d_n(y_{n+2}) &= d(y_n, y_{n+1}, y_{n+2}) = d(Tx_n, Tx_{n+1}, y_n) \\ &\leq a \max \{d(fx_n, fx_{n+1}, y_n), d(fx_n, Tx_n, y_n), d(fx_{n+1}, Tx_{n+1}, y_n)\} \\ &\quad, \frac{1}{2} [M(x_n, x_{n+1}, y_n) + m(x_n, x_{n+1}, y_n)] \} + c [M(x_n, x_{n+1}, y_n) + h m(x_n, x_{n+1}, y_n)] \\ &= a \max \{d(y_n, y_{n+1}, y_n), d(y_n, y_{n+1}, y_n), d(y_{n+1}, y_{n+2}, y_n)\} \\ &\quad, \frac{1}{2} [M(x_n, x_{n+1}, y_n) + m(x_n, x_{n+1}, y_n)] \} + c [M(x_n, x_{n+1}, y_n) + h m(x_n, x_{n+1}, y_n)] \end{aligned}$$

where a and c are evaluated at (x_n, x_{n+1}) .

Since

$$\begin{aligned} m(x_n, x_{n+1}, y_n) &= \min \{d(fx_n, Tx_{n+1}, y_n), d(fx_{n+1}, Tx_n, y_n)\} \\ &= \min \{d(y_n, y_{n+2}, y_n), d(y_{n+1}, y_{n+1}, y_n)\} = 0 \end{aligned}$$

And

$$\begin{aligned} M(x_n, x_{n+1}, y_n) &= \max\{d(fx_n, Tx_{n+1}, y_n), d(fx_{n+1}, Tx_n, y_n)\} \\ &= \max\{d(y_n, y_{n+2}, y_n), d(y_{n+1}, y_{n+1}, y_n)\} = 0. \end{aligned}$$

Hence from (2.1), we have

$$d_n(y_{n+2}) \leq a d_n(y_{n+2})$$

From (1.6), we have

$$(2.2) \quad d_n(y_{n+2}) = 0.$$

We shall prove that $\{d_n(u)\}_{n \in \mathbb{N} \cup \{0\}}$ is a non-increasing sequence in \mathbb{R}^+ . For all $u \in X$, on the contrary, assume that $d_{n+1}(u) > d_n(u)$. Again applying (1.5), we have

$$\begin{aligned} (2.3) \quad d_{n+1}(u) &= d(y_{n+1}, y_{n+2}, u) = d(Tx_n, Tx_{n+1}, u) \\ &\leq a \max\{d(fx_n, fx_{n+1}, u), d(fx_n, Tx_n, u), d(fx_{n+1}, Tx_{n+1}, u) \\ &\quad, \frac{1}{2}[M(x_n, x_{n+1}, u) + m(x_n, x_{n+1}, u)]\} + c[M(x_n, x_{n+1}, u) + h m(x_n, x_{n+1}, u)] \\ &= a \max\{d_n(u), d_n(u), d_{n+1}(u), \frac{1}{2}[M(x_n, x_{n+1}, u) + m(x_n, x_{n+1}, u)]\} \\ &\quad + c[M(x_n, x_{n+1}, u) + h m(x_n, x_{n+1}, u)] \end{aligned}$$

where a and c are evaluated at (x_n, x_{n+1}) .

Since

$$\begin{aligned} m(x_n, x_{n+1}, u) &= \min\{d(fx_n, Tx_{n+1}, u), d(fx_{n+1}, Tx_n, u)\} \\ &= \min\{d(y_n, y_{n+2}, u), d(y_{n+1}, y_{n+1}, u)\} = 0 \\ M(x_n, x_{n+1}, u) &= \max\{d(fx_n, Tx_{n+1}, u), d(fx_{n+1}, Tx_n, u)\} \\ &= \max\{d(y_n, y_{n+2}, u), d(y_{n+1}, y_{n+1}, u)\} \\ &= d(y_n, y_{n+2}, u). \end{aligned}$$

Then from (2.2), we have

$$\begin{aligned} M(x_{n+1}, x_{n+2}, u) &= d(y_n, y_{n+2}, u) \\ &\leq d(y_n, y_{n+2}, y_{n+1}) + d(y_n, y_{n+1}, u) + d(y_{n+1}, y_{n+2}, u) \\ &\leq 2d_{n+1}(u) \end{aligned}$$

The inequality (2.3) gives

$$d_{n+1}(u) \leq (a + 2c) d_{n+1}(u)$$

a contradiction. Thus, our supposition that $d_{n+1}(u) > d_n(u)$ was wrong. Therefore $\{d_n(u)\}_{n \in \mathbb{N} \cup \{0\}}$ is a non-increasing sequence of non-negative real numbers. Therefore, for all n we have

$$(2.4) \quad d_{n+1}(u) \leq d_n(u).$$

Now, we shall prove that for any $n, m \in \mathbb{N}$, $d_n(y_m) = 0$.

Let $n, m \in \mathbb{N}$ and if $n \geq m$ and $u = y_m$, then from (2.4), we have

$$(2.5) \quad d_n(y_m) \leq d_{n-1}(y_m) \leq \dots \leq d_m(y_m) = 0.$$

If $n < m$, then from (2.5), we have

$$\begin{aligned} d_n(y_m) &= d(y_n, y_{n+1}, y_m) \\ &\leq d(y_n, y_{n+1}, y_{m-1}) + d(y_n, y_{m-1}, y_m) + d(y_{m-1}, y_{n+1}, y_m) \\ &= d_n(y_{m-1}) + d_{m-1}(y_n) + d_{m-1}(y_{n+1}) \\ &= d_n(y_{m-1}) \leq d_n(y_{m-2}) \leq \dots \leq d_n(y_{n+1}) = 0. \end{aligned}$$

Thus for any $n, m \in \mathbb{N}$,

$$(2.6) \quad d_n(y_m) = 0$$

Next we shall prove that $d(y_i, y_j, y_k) = 0$ for all $i, j, k \in \mathbb{N}$. Without loss of generality, we may assume that $i \leq j$, it follows that

$$\begin{aligned} d(y_i, y_j, y_k) &\leq d(y_i, y_j, y_{i+1}) + d(y_i, y_{i+1}, y_k) + d(y_{i+1}, y_j, y_k) \\ &= d_i(y_j) + d_i(y_k) + d(y_{i+1}, y_j, y_k) \end{aligned}$$

$$= d(y_{i+1}, y_j, y_k)$$

Similarly,

$$d(y_{i+1}, y_j, y_k) \leq d(y_{i+2}, y_j, y_k)$$

Inductively, we have

$$(2.7) \quad d(y_i, y_j, y_k) \leq d(y_{j-1}, y_j, y_k) = d_j(y_k) = 0$$

We claim that $\lim_{n \rightarrow \infty} d_n(u) = 0$.

Applying (1.5), we have

$$\begin{aligned} (2.8) \quad d(y_{n-1}, Tx_n, u) &= d(Tx_{n-2}, Tx_n, u) \\ &\leq a \max\{d(fx_{n-2}, fx_n, u), d(fx_{n-2}, Tx_{n-2}, u), d(fx_n, Tx_n, u) \\ &\quad, \frac{1}{2} [M(x_{n-2}, x_n, u) + m(x_{n-2}, x_n, u)]\} + c[M(x_{n-2}, x_n, u) + hm(x_{n-2}, x_n, u)] \\ &= a \max\{d(y_{n-2}, y_n, u), d(y_{n-2}, y_{n-1}, u), d(y_n, y_{n+1}, u) \\ &\quad, \frac{1}{2} [M(x_{n-2}, x_n, u) + m(x_{n-2}, x_n, u)]\} + c[M(x_{n-2}, x_n, u) + hm(x_{n-2}, x_n, u)] \end{aligned}$$

where a and c are evaluated at (x_{n-2}, x_n) .

Since by using (2.4), (2.5) and (2.6), we have

$$\begin{aligned} d(y_{n-2}, y_n, u) &\leq d(y_{n-2}, y_n, y_{n-1}) + d(y_{n-2}, y_{n-1}, u) + d(y_{n-1}, y_n, u) \\ &\leq d_{n-2}(y_n) + d_{n-2}(u) + d_{n-1}(u) \\ &\leq 2d_{n-2}(u) \\ d(y_{n-2}, y_{n+1}, u) &\leq d(y_{n-2}, y_{n+1}, y_{n-1}) + d(y_{n-2}, y_{n-1}, u) + d(y_{n-1}, y_{n+1}, u) \\ &\leq d(y_{n-2}, y_{n+1}, y_{n-1}) + d_{n-2}(u) + d(y_{n-1}, y_{n+1}, y_n) + d(y_{n-1}, y_n, u) \\ &\quad + d(y_n, y_{n+1}, u) \\ &= d_{n-2}(u) + d_{n-1}(u) + d_n(u) \end{aligned}$$

$$\leq 3d_{n-2}(u)$$

$$m(x_{n-2}, x_n, u) = \min\{d(fx_{n-2}, Tx_n, u), d(fx_n, Tx_{n-2}, u)\}$$

$$= \min\{d(y_{n-2}, y_{n+1}, u), d(y_n, y_{n-1}, u)\}$$

$$\leq \min\{3d_{n-2}(u), d_{n-1}(u)\}$$

$$= d_{n-1}(u)$$

$$M(x_{n-2}, x_n, u) = \max\{d(fx_{n-2}, Tx_n, u), d(fx_n, Tx_{n-2}, u)\}$$

$$= \max\{d(y_{n-2}, y_{n+1}, u), d(y_n, y_{n-1}, u)\}$$

$$\leq \max\{3d_{n-2}(u), d_{n-1}(u)\} = 3d_{n-2}(u)$$

Using above all inequalities and (1.7), the inequality (2.8) gives

$$\begin{aligned} (2.9) \quad d(y_{n-1}, y_{n+1}, u) &\leq a \max\left\{2d_{n-2}(u), d_{n-2}(u), d_n(u), \frac{1}{2}[3d_{n-2}(u) + d_{n-1}(u)]\right\} \\ &\quad + c[3d_{n-2}(u) + hd_{n-1}(u)] \\ &\leq [2a + c(3 + h)]d_{n-2}(u) \\ &= [2 - c(1 - h)]d_{n-2}(u) \end{aligned}$$

Again from (1.5), we have

$$\begin{aligned} (2.10) \quad d(y_n, y_{n+1}, u) &= d(Tx_{n-1}, Tx_n, u) \\ &\leq a \max\{d(fx_{n-1}, fx_n, u), d(fx_{n-1}, Tx_{n-1}, u), d(fx_n, Tx_n, u) \\ &\quad , \frac{1}{2}[M(x_{n-1}, x_n, u) + m(x_{n-1}, x_n, u)]\} + c[M(x_{n-1}, x_n, u) + hm(x_{n-1}, x_n, u)] \\ &= a \max\{d(y_{n-1}, y_n, u), d(y_{n-1}, y_n, u), d(y_n, y_{n+1}, u) \\ &\quad , \frac{1}{2}[M(x_{n-1}, x_n, u) + m(x_{n-1}, x_n, u)]\} + c[M(x_{n-1}, x_n, u) + hm(x_{n-1}, x_n, u)] \end{aligned}$$

where a and c are evaluated at (x_{n-1}, x_n) .

Since

$$\begin{aligned} m(x_{n-1}, x_n, u) &= \min\{d(fx_{n-1}, Tx_n, u), d(fx_n, Tx_{n-1}, u)\} \\ &= \min\{d(y_{n-1}, y_{n+1}, u), d(y_n, y_n, u)\} = 0 \\ M(x_{n-2}, x_n, u) &= \max\{d(fx_{n-1}, Tx_n, u), d(fx_n, Tx_{n-1}, u)\} \\ &= \max\{d(y_{n-1}, y_{n+1}, u), d(y_n, y_n, u)\} \\ &= d(y_{n-1}, y_{n+1}, u) \end{aligned}$$

Using (2.4) and (2.9), the inequality (2.10) gives

$$\begin{aligned} d_n(u) &\leq \alpha d_{n-1}(u) + c d(y_{n-1}, y_{n+1}, u) \\ &\leq \alpha d_{n-2}(u) + c [2 - c(1 - h)] d_{n-2}(u) \\ &= [1 - c^2(1 - h)] d_{n-2}(u) \end{aligned}$$

Hence

$$d_n(u) \leq [1 - \beta^2(1 - h)] d_{n-2}(u)$$

Proceeding in this manner, we obtain

$$(2.11) \quad d_n(u) \leq (1 - \beta^2(1 - h))^{\left[\frac{n}{2}\right]} d_0(u)$$

where $\left[\frac{n}{2}\right]$ stands for the greatest integer not exceeding $\frac{n}{2}$. Since $\beta = \inf\{c(x, y) : x, y \in X\} > 0$ and $h \in (0, 1)$, which implies that

$$(2.12) \quad \lim_{n \rightarrow \infty} d_n(u) = 0.$$

Now, we prove that $\{y_n\}$ is Cauchy. Suppose to the contrary, that $\{y_n\}$ is not a Cauchy sequence in X . Then for every $\epsilon > 0$, there exists $u \in X$ and strictly increasing sequences $\{m_k\}, \{n_k\}$ of positive integers such that $m_k > n_k \geq k$ with

$$(2.13) \quad d(y_{m_k}, y_{n_k}, u) \geq \epsilon$$

Without loss of generality, we can suppose that also

$$(2.14) \quad m_k > n_k \geq k, \quad d(y_{m_k}, y_{n_k}, u) \geq \epsilon, \quad d(y_{n_k}, y_{m_{k-2}}, u) < \epsilon$$

From (2.14) and the tetrahedral inequality (that holds for a 2-metric space), we have

$$\begin{aligned}
 (2.15) \quad \epsilon &\leq d(y_{m_k}, y_{n_k}, u) \\
 &\leq d(y_{m_k}, y_{n_k}, y_{m_k-2}) + d(y_{m_k}, y_{m_k-2}, u) + d(y_{m_k-2}, y_{n_k}, u) \\
 &\leq d(y_{m_k-2}, y_{n_k}, u) + d(y_{m_k}, y_{n_k}, y_{m_k-2}) + d(y_{m_k-1}, y_{m_k-2}, u) \\
 &\quad + d(y_{m_k}, y_{m_k-1}, u) + d(y_{m_k}, y_{m_k-2}, y_{m_k-1}) \\
 &\leq \epsilon + d(y_{m_k}, y_{n_k}, y_{m_k-2}) + d_{m_k-2}(u) + d_{m_k-1}(u) + d_{m_k-2}(y_{m_k})
 \end{aligned}$$

On letting $k \rightarrow +\infty$ in (2.15) and using (2.2), (2.7), (2.12), we get

$$(2.16) \quad \lim_{k \rightarrow +\infty} d(y_{m_k}, y_{n_k}, u) = \epsilon$$

It follows from (2.14) that

$$\begin{aligned}
 0 &< d(y_{n_k}, y_{m_k}, u) - d(y_{n_k}, y_{m_k-2}, u) \\
 &\leq d(y_{n_k}, y_{m_k-2}, u) + d(y_{m_k-2}, y_{m_k}, u) + d(y_{n_k}, y_{m_k}, y_{m_k-2}) - d(y_{n_k}, y_{m_k-2}, u) \\
 &= d(y_{m_k-2}, y_{m_k}, u) + d(y_{n_k}, y_{m_k}, y_{m_k-2}) \\
 &\leq d_{m_k-2}(y_{m_k}) + d_{m_k-2}(u) + d_{m_k-1}(u) + d(y_{n_k}, y_{m_k}, y_{m_k-2})
 \end{aligned}$$

On making $k \rightarrow +\infty$, we immediately obtain that:

$$(2.17) \quad \lim_{k \rightarrow +\infty} d(y_{n_k}, y_{m_k-2}, u) = \epsilon$$

Note that

$$\begin{aligned}
 |d(y_{n_k}, y_{m_k-1}, u) - d(y_{n_k}, y_{m_k}, u)| &\leq d_{m_k-1}(u) + d_{m_k-1}(y_{n_k}) \\
 |d(y_{n_k+1}, y_{m_k}, u) - d(y_{n_k}, y_{m_k}, u)| &\leq d_{n_k}(u) + d_{n_k}(y_{m_k}) \\
 |d(y_{n_k+1}, y_{m_k-1}, u) - d(y_{n_k}, y_{m_k}, u)| &\leq d_{n_k}(u) + d_{m_k-1}(y_{n_k}) \\
 &\quad + d_{m_k-1}(u) + d_{n_k}(y_{m_k-1})
 \end{aligned}$$

On letting $k \rightarrow +\infty$, in these inequalities and by using inequalities (2.2), (2.7), (2.12), (2.15) and (2.17), we obtain;

$$(2.18) \quad \lim_{k \rightarrow +\infty} d(y_{n_k}, y_{m_k-1}, u) = \epsilon, \quad \lim_{k \rightarrow +\infty} d(y_{n_k+1}, y_{m_k}, u) = \epsilon,$$

$$\lim_{k \rightarrow +\infty} d(y_{n_k+1}, y_{m_k-1}, u) = \epsilon.$$

Now, using (1.5), we have

$$(2.19) \quad d(y_{m_k}, y_{n_k+1}, u) = d(Tx_{m_k-1}, Tx_{n_k}, u)$$

$$\leq a \max\{d(fx_{m_k-1}, fx_{n_k}, u), d(fx_{m_k-1}, Tx_{m_k-1}, u), d(fx_{n_k}, Tx_{n_k}, u)$$

$$, \frac{1}{2} [M(x_{m_k-1}, x_{n_k}, u) + m(x_{m_k-1}, x_{n_k}, u)]\}$$

$$+ c [M(x_{m_k-1}, x_{n_k}, u) + h m(x_{m_k-1}, x_{n_k}, u)]$$

$$= a \max\{d(y_{m_k-1}, y_{n_k}, u), d(y_{m_k-1}, y_{m_k}, u), d(y_{n_k}, y_{n_k+1}, u)$$

$$, \frac{1}{2} [M(x_{m_k-1}, x_{n_k}, u) + m(x_{m_k-1}, x_{n_k}, u)]\}$$

$$+ c [M(x_{m_k-1}, x_{n_k}, u) + h m(x_{m_k-1}, x_{n_k}, u)]$$

where a and c are evaluated at (x_{m_k-1}, x_{n_k}) and

$$m(x_{m_k-1}, x_{n_k}, u) = \min\{d(fx_{m_k-1}, Tx_{n_k}, u), d(fx_{n_k}, Tx_{m_k-1}, u)\}$$

$$= \min\{d(y_{m_k-1}, y_{m_k+1}, u), d(y_{n_k}, y_{m_k}, u)\}$$

$$M(x_{m_k-1}, x_{n_k}, u) = \max\{d(fx_{m_k-1}, Tx_{n_k}, u), d(fx_{n_k}, Tx_{m_k-1}, u)\}$$

$$= \max\{d(y_{m_k-1}, y_{m_k+1}, u), d(y_{n_k}, y_{m_k}, u)\}$$

On letting $k \rightarrow +\infty$ in (2.19), using (2.12), (2.16) and (2.18), we have

$$\epsilon \leq a \max\{\epsilon, 0, 0, \frac{1}{2} [\epsilon + \epsilon]\} + c[\epsilon + h\epsilon]$$

$$= (a + c(1 + h))\epsilon$$

$$= (1 - c(1 - h))\varepsilon$$

This is a contradiction, since $\beta = \inf\{c(x, y) : x, y \in X\} > 0$ and $h \in (0, 1)$. Thus, our supposition was wrong and therefore, $\{y_n\}$ is a Cauchy Sequence in X .

For cases (a) and (b) suppose that X is complete. Then Cauchy sequence $\{y_n\}$ will converge to a point p in X and then $fx_n \rightarrow p$ and $Tx_n \rightarrow p$ as $n \rightarrow +\infty$.

Case (a): Suppose that f is surjective. Then there exists a point z in X such that $p = fz$.

From (1.5), we have

$$\begin{aligned} (2.20) \quad d(fz, Tz, u) &\leq d(fz, Tz, y_{n+1}) + d(fz, y_{n+1}, u) + d(y_{n+1}, Tz, u) \\ &= d(fz, y_{n+1}, u) + d(fz, Tz, y_{n+1}) + d(Tx_n, Tz, u) \\ &\leq d(fz, y_{n+1}, u) + d(fz, Tz, y_{n+1}) \\ &\quad + a \max \left\{ d(fx_n, fz, u), d(fx_n, Tx_n, u), d(fz, Tz, u), \frac{1}{2} [M(x_n, z, u) + m(x_n, z, u)] \right\} \\ &\quad + c [M(x_n, z, u) + h m(x_n, z, u)] \end{aligned}$$

Since

$$\begin{aligned} \lim_{n \rightarrow +\infty} M(x_n, z, u) &= \lim_{n \rightarrow +\infty} \max \{ d(fx_n, Tz, u), d(fz, Tx_n, u) \} \\ &= d(fz, Tz, u) \end{aligned}$$

and

$$\begin{aligned} \lim_{n \rightarrow +\infty} m(x_n, z, u) &= \lim_{n \rightarrow +\infty} \min \{ d(fx_n, Tz, u), d(fz, Tx_n, u) \} \\ &= 0. \end{aligned}$$

Taking limit $n \rightarrow +\infty$ in the inequality (2.20), we have

$$d(fz, Tz, u) \leq \sup_{x, y \in X} (a + c) d(fz, Tz, u) < d(fz, Tz, u)$$

implies that $fz = Tz = p$.

Case (b): Suppose f is continuous. Then since $\lim_{n \rightarrow +\infty} y_n = p$, we have $\lim_{n \rightarrow +\infty} f y_n = f p$ and then $\lim_{n \rightarrow +\infty} f f x_n = f p$. Also f and T are compatible and $\lim_{n \rightarrow +\infty} f x_n = \lim_{n \rightarrow +\infty} T x_n = \lim_{n \rightarrow +\infty} y_{n+1} = p$. Hence

$$(2.21) \quad \lim_{n \rightarrow +\infty} d(f T x_n, T f x_n, u) = 0$$

Note that

$$\begin{aligned} d(f f x_n, T f x_n, u) &\leq d(f f x_n, T f x_n, f T x_n) + d(f f x_n, f T x_n, u) + d(f T x_n, T f x_n, u) \\ &= d(f f x_n, T f x_n, f T x_n) + d(f f x_n, f f x_{n+1}, u) + d(f T x_n, T f x_n, u) \end{aligned}$$

On taking limit $n \rightarrow +\infty$ and using (2.21), we have $\lim_{n \rightarrow +\infty} d(f f x_n, T f x_n, u) = 0$. Since $\lim_{n \rightarrow +\infty} f f x_n = f p$, it follows that $\lim_{n \rightarrow +\infty} T f x_n = f p$.

Applying (1.5) again, we have

$$\begin{aligned} (2.22) \quad d(f p, T p, u) &\leq d(f p, T p, f y_{n+1}) + d(f p, f y_{n+1}, u) + d(f y_{n+1}, T p, u) \\ &\leq d(f p, T p, f y_{n+1}) + d(f p, f y_{n+1}, u) + d(T p, f T x_n, u) \\ &\leq d(f p, T p, f y_{n+1}) + d(f p, f y_{n+1}, u) + d(T p, f T x_n, T f x_n) \\ &\quad + d(T p, T f x_n, u) + d(T f x_n, f T x_n, u) \\ &\leq d(f p, T p, f y_{n+1}) + d(f p, f y_{n+1}, u) + d(T p, f T x_n, T f x_n) \\ &\quad + d(T f x_n, f T x_n, u) + a \max\{d(f p, f f x_n, u), d(f p, T p, u), d(f f x_n, T f x_n, u) \\ &\quad, \frac{1}{2}[M(p, f x_n, u) + m(p, f x_n, u)]\} + c[M(p, f x_n, u) + h m(p, f x_n, u)] \end{aligned}$$

Note that

$$\begin{aligned} \lim_{n \rightarrow +\infty} M(p, f x_n, u) &= \lim_{n \rightarrow +\infty} \max\{d(f p, T f x_n, u), (f f x_n, T p, u)\} \\ &= \lim_{n \rightarrow +\infty} \max\{d(f p, T f x_n, u), (f f x_n, T p, u)\} \\ &= d(f p, T p, u) \end{aligned}$$

and

$$\begin{aligned}\lim_{n \rightarrow +\infty} m(p, f x_n, u) &= \lim_{n \rightarrow +\infty} \min\{d(fp, T f x_n, u), (f f x_n, T p, u)\} \\ &= \lim_{n \rightarrow +\infty} \min\{d(fp, T f x_n, u), (f f x_n, T p, u)\} \\ &= 0.\end{aligned}$$

On letting $n \rightarrow +\infty$ in the inequality (2.22), we have

$$(2.23) \quad d(fp, Tp, u) \leq \sup_{x, y \in X} (a + c) d(fp, Tp, u)$$

implies that $fp = Tp$.

Case (c): In this case $p \in f(X)$. Let $z \in f^{-1}(p)$. Then $p = fz$ and the proof is complete by case (a).

Case (d): In this case $p \in T(X) \subseteq f(X)$ and the proof is complete by case (c).

Finally, we shall prove that f and T have at most one coincidence point. On the contrary, suppose that f and T have two coincidence points p and q . Then from (1.5) with a and b evaluated at (p, q) , we have

$$\begin{aligned}(2.24) \quad d(Tp, Tq, u) &\leq a \max\{d(fp, fq, u), d(fp, Tp, u), d(fq, Tq, u), \frac{1}{2}[M(p, q, u) + m(p, q, u)]\} \\ &\quad + c[M(p, q, u) + h m(p, q, u)] \\ &= [a + c(1 + h)] d(Tp, Tq, u)\end{aligned}$$

because

$$\begin{aligned}M(p, q, u) &= \max\{d(fp, Tq, u), d(fq, Tp, u)\} \\ &= d(Tq, Tp, u) \\ m(p, q, u) &= \min\{d(fp, Tq, u), d(fq, Tp, u)\} \\ &= d(Tq, Tp, u)\end{aligned}$$

Hence by (1.7),

$$(2.25) \quad d(Tp, Tq, u) \leq [1 - c(1 - h)] d(Tp, Tq, u)$$

implying $Tp = Tq$ by (1.6) and hence $fp = fq$.

Corollary 2.2 Let (X, d) be a complete 2-metric space and T is self mapping of X satisfying (1.5) with $f = I$, the identity map on X , where $h = 1, a$ and b satisfying (1.6) and (1.7). Then T has a unique fixed point and at this fixed point T is continuous.

Proof The existence and uniqueness of the fixed point comes from Theorem 2.1 by setting $f = I$. To prove continuity, let $\{y_n\} \subset X$ with $\lim_{n \rightarrow +\infty} y_n = p$, p the unique fixed point of T .

We apply inequality (1.6), where a, c are evaluating at (y_n, p) .

$$\begin{aligned}
 (2.26) \quad d(Ty_n, Tp, u) &\leq a \max \left\{ d(y_n, p, u), d(y_n, Ty_n, u), d(p, Tp, u), \frac{1}{2} [M(y_n, p, u) + m(y_n, p, u)] \right\} \\
 &\quad + c [M(y_n, p, u) + m(y_n, p, u)] \\
 &\leq a (d(y_n, p, u) + d(p, Ty_n, u)) + c [d(p, Ty_n, u) + d(y_n, p, u)] \\
 &= (a + c) d(y_n, p, u) + (a + c) d(p, Ty_n, u) \\
 &= (1 - c) d(y_n, p, u) + (1 - c) d(p, Ty_n, u)
 \end{aligned}$$

Hence

$$(2.27) \quad d(Ty_n, Tp, u) \leq (1 - \beta) d(y_n, p, u) + (1 - \beta) d(p, Ty_n, u)$$

Since $\beta = \inf \{c(x, y) : x, y \in X\} > 0$. Hence we get

$$(2.28) \quad d(Ty_n, Tp, u) \leq \left(\frac{1}{\beta} - 1\right) d(y_n, p, u)$$

Taking the limit as $n \rightarrow +\infty$ yields

$$\lim_{n \rightarrow +\infty} Ty_n = Tp.$$

Therefore T is continuous at p .

CONFLICT OF INTEREST

No conflict of interest was declared by the authors.

AUTHOR'S CONTRIBUTIONS

All authors contributed equally and significantly to writing this paper. All authors read and approved the final manuscript.

5. REFERENCES

- [1] Banach, S., Sur les operations dans les ensembles abstraits et leur application aux equations integrales, Fund. Math., 3(1922), 133-181.
- [2] Belluce, L. P., Kirk, W. A., Nonexpansive mappings and fixed points in Banach space. 111. J. Math. 11 (1967), 474-479.
- [3] Bogin, J., A generalization of a fixed point theorem of Gebel, Kirk and Shimi, Canad. Math. Bull., 19(1976), 7-12.
- [4] Browder, F. E., Nonexpansive nonlinear operators in a Banach space. Proc. Nat. Acad. Sci. U.S.A. 54 (1965), 1041-1044.
- [5] Chandra, M., Mishra, S., Singh, S., Rhoades, B.E., Coincidence and fixed points of nonexpansive type multi-valued and single-valued maps, Indian J. Pure Appl. Math., 26 (5):393-401, 1995.
- [6] Ćirić, Lj. B., A new class of nonexpansive type mappings and fixed points, Czechoslovak Mathematical Journal, Vol. 49 (1999), No. 4, 891—899.
- [7] Ćirić, Lj. B., On some non-expansive type mappings and fixed points, Indian J. Pure Appl. Math., 24(3):145-149, 1993.
- [8] Gähler, S., 2-metrische Räume und ihre topologische structure, Math Nachr, 26(1963), 115-148.
- [9] Gähler, S., Über die uniformisierbarkeit 2-metrische Räume, Math Nachr, 28(1965), 235-244.
- [10] Gähler, S., Zur geometrie 2-metrische Räume, Rev Raum Math Pures Et Appl., 11(1966), 655-664.
- [11] Gregus, M., A fixed point theorem in Banach spaces, Boll. Un. Mat. Ital.. A, 5 (1980), 193-198.
- [12] Iseki, K., Fixed point theorems in 2-metric spaces, Math Sem Notes, 3(1975), 133-136.
- [13] Iseki, K., On common fixed point theorems of mappings. Proc. Japan Acad. Ser. A Math. Sci. 50 (1974), 408—409.
- [14] Jhade, P.K., Saluja, A. S., Kushwah, R., Coincidence & Fixed Points of Nonexpansive Type Multi-Valued & Single Valued Maps, European Journal of Pure And Applied Mathematics Vol. 4, No. 4, 2011, 330-339.
- [15] Liu, Z. Zhang, F., Characterizations of common fixed points in 2-metric spaces, Rostock. Math. Kolloq., 55(2001), 49-64.
- [16] Murthy, P.P., Chang, S.S., Cho, Y.J., Sharma, B.K., Compatible mappings of type (A) and common fixed point theorem, Kyungpook Math J, 32(1992), 203-216.
- [17] Pant R. P., Common fixed points of four mappings. Bull. Calcutta Math. Soc., 90(4) (1998), 281-286.
- [18] Pathak H. K., Kang S. M., Baek J. H., Weakly compatible mappings of type (A) and common fixed points, Kyungpook Math. J., 35(2) (1995), 345-359.
- [19] Rhoades, B. E., A generalization of a fixed point theorem of Bogin. Math. Sem. Notes, Kobe Univ. 6 (1978), 1—7.
- [20] Singh, S.L., Adiga, C., GiniSwami, A fixed point theorem in 2-metric space and an application, Journal of Natural & Physical Science, 15(1-2)(2001), 55-64.
- [21] Ughade, M., Daheriya, R. D., Using Implicit Relation to Prove Unified Fixed Point Theorems in 2-Metric Spaces, International J. of Math. Sci. & Engg. Appls. Vol. 9 No. IV (2015), pp. 57-80.