

THERMO-ACOUSTIC MOLECULAR INTERACTION STUDIES IN BINARY LIQUID MIXTURES OF PROPYL AMINE AND BENZENE USING ULTRASONIC TECHNIQUE AT 301K.

Ruman Singh¹, K.Y. Singh² and R.C. Verma*

¹ Deptt. of Physics, Sun Rise University Alwar (Raj.) India

² Deptt. of Physics, B.S.A. College, Mathura (U.P.) India

*Deptt. of Chemistry, Janta College, Bakewar (U.P.) India

Email : jcb.rajesh@gmail.com

ABSTRACT

The ultrasonic studies in liquids are great use in understanding the nature and strength of molecular interaction. The thermo-acoustical parameters for binary liquid mixtures of propyl amine and benzene have been estimated from the measured values of ultrasonic velocity (v), density (ρ) and viscosity (η). Using the measured data, some of acoustic parameters such as isentropic compressibility (β_s) and intermolecular free length (L_f) are evaluated at the temperature 303K. The present paper represents the nonlinear variation of ultrasonic velocity and thermo-acoustical parameters lead to dipole-induced dipole interaction between propyl amine and benzene molecules. The behavior of these parameters with composition of the mixture has been discussed in terms of molecular interaction between the components of the liquids.

Keyword: *ultrasonic velocity, acoustical parameters, molecular interaction, propyl amine, benzene.*

INTRODUCTION

Ultrasonic study is very much useful for characterizing the physic-chemical behavior of liquid mixtures and measurements are used to study molecular interactions in liquids.¹⁻² The method of studying in molecular interaction from the knowledge of variation of acoustic parameters along with their excess values with change in mole fraction gives an insight into the molecular process.³⁻⁴ The increase or decrease in ultrasonic velocities have been employed in understanding the nature of molecular interaction in the pure liquid binary mixtures.⁵⁻¹⁰ The study of liquid mixtures containing of polar and non-polar components find applications in industrial and technological process.¹¹

The mixing of different give rise to solutions that generally do not behave ideally.¹²⁻¹³ Further those properties have been widely used to study the molecular interaction between the various species in the mixture.¹⁴⁻¹⁵

In the present study ultrasonic velocity, density and viscosity were measured experimentally for binary system namely propyl amine + benzene at 301K. From the measured data , thermo-acoustical parameters have been computed and the results are analysed in the light of molecular interaction.

MATERIALS AND METHODS

Propyl amine and benzene were used after single distillation. Binary mixtures were prepared by mixing known volume of each liquid in air tight Stoppard glass bottle. Care was taken to avoid contamination during mixing.

Ultrasonic velocity was measured by Ultrasonic Interferometer M-80 manufactured by M/S Mittal Enterprises, New Delhi having accuracy of about $\pm 0.057\%$.

Density of pure liquid and binary mixtures was measured by using double walled Picknometer. The Picknometer was calibrated with distilled water. The value obtained were tally with the literature values. The viscosities have been determined by using Ostwald viscometer. The accuracy in viscosity measurement was ± 0.0002 c.p.

Isentropic compressibility (β_s) has been calculated from ultrasonic velocity (v) and the density (ρ) usin the equation as:

$$\beta_s = 1/v^2\rho \quad (1)$$

Intermolecular free length (L_f) has been determined as:

$$L_f = KT(\beta_s)^{1/2} \quad (2)$$

Where KT is a Jacobson's constant.

Table-1: Experimental values of ultrasonic velocity (v), density (ρ) and viscosity (η) of pure liquids at 303K.

Liquid	Ultrasonic Velocity	Density	Viscosity
Propyl amine	1160	0.6980	0.3780
Benzene	1294	0.8690	0.5610

Table-2: : Experimental values of ultrasonic velocity (v), density (ρ) and viscosity (η) for the binary liquid mixture of propyl amine and benzene at 301K.

Mole Fraction of propyl amine (X1)	Ultrasonic Velocity (v) ms^{-1}	Density (ρ) Gm.ml^{-1}	Viscosity (η) cp
0.0000	1294	0.8690	0.5610
0.1316	1276	0.8448	0.5381
0.2613	1258	0.8215	0.5152
0.3890	1241	0.7988	0.4924
0.5148	1224	0.7765	0.4697
0.6388	1208	0.7556	0.4469
0.7610	1192	0.7359	0.4240
0.8813	1176	0.7170	0.4011
1.0000	1160	0.6980	0.3780

Table-3: Experimental values of isentropic compressibility (β_s) and intermolecular free length (L_f) for the binary liquid mixture of propyl amine and benzene at 301K

Mole Fraction of methyl amine X_1	Isentropic Compressibility(β) $\text{Cm}^2\text{dyne}^{-1}\times 10^{12}$	Intermolecular Free length (L_f) \AA
0.0000	68.72	0.5205
0.1316	72.70	0.5372
0.2613	76.91	0.55.37
0.3890	81.28	0.5700
0.5148	85.95	0.5861
0.6388	90.69	0.60.18
0.7610	95.63	0.61.74
0.8813	100.84	0.63.22
1.0000	106.47	0.64.79

Graphs: Fig. 1-5 show variation of ultrasonic velocity (v), density (ρ), viscosity (η), isentropic compressibility(β_s) and intermolecular free length (L_f) with respect to mole fraction at temperature 301K

Fig.1: Variation of ultrasonic velocity with mole fraction.




Fig.2: Variation of density with mole fraction.




Fig.3: Variation of viscosity with mole fraction




Fig.4: Variation of isentropic compressibility with mole fraction.





Fig.5: Variation of intermolecular free length with mole fraction.



RESULTS AND DISCUSSION

The experimentally measured values of ultrasonic velocity, density and viscosity for pure liquids at 301K are presented in Table-1. Experimental values of ultrasonic velocity, density and viscosity for binary mixture at 301K are given in Table-2. The thermodynamic parameters such as isentropic compressibility (β_s) and intermolecular free length (L_f) are listed in Table-3. The variation of

ultrasonic velocity, density and viscosity at 298K are shown in Fig.1,2 and 3 respectively. While other thermodynamic parameters such as isentropic compressibility (β_s) and intermolecular free length (L_f) at 298K are shown in Fig. 4 and 5 respectively.

From Table-2 it is observed that, the density (ρ) decreases with increase in mole fraction for propyl amine and benzene system while viscosity (η) decreases with increase in mole fraction and ultrasonic velocity (v) decreases with increasing mole fraction. The decrease in ultrasonic velocity are due to the increase in isentropic compressibility and intermolecular free length of the liquid mixtures. This may lead to presence of dispersive force (London force) between the molecules of the liquid mixture.¹⁶ The isentropic compressibility and intermolecular free length are the deciding factors of ultrasonic velocity in binary mixtures.

As benzene is non-polar molecule does not possess dipole moment, when it interacts with propyl amine which is polar molecule possess dipole moment then benzene possess induced dipole moment. This induced dipole-dipole interaction between benzene and propyl amine molecules.

CONCLUSION

From ultrasonic velocity, related acoustic parameters for methyl amine with o-xylene for various concentration at 298K, it has been found that there exists a dipole-induced dipole interaction between methyl amine and o-xylene.

REFERENCES:

1. Kannappam,V. and Chidambara Vinayagam,S.(2006), Ultrasonic study of ion-solvent interaction in aqueous and nonaqueous solutions of transition and inner transition metal ions.,*Ind.J.Pure & Appl.Phys.*,44(9),670-676.
2. Voleisiene,B.and Voleisis,A.,(2008), Ultrasound velocity measurements in liquid media.*Ultragarsas (Ultrasound)*,63(4),7-19.
3. Jain,D.V.S. *properties* and Dhar,N.S.(1992), *Excess molar enthalpies of (Benzene or methyl benzene or ethyl benzene + 2-methyl ethyl benzene) at the temperature 298.15K, 308.15K and 318.15..J.Chem.Thermodynamics.*,24(10),1027-1031..

4. Largemann,R.T. and Dumbard,W.S.(1992) , *Relationship between the velocity of sound and other properties of liquids* *J.Phys.Chem.*,**1945**,49(5),428-436.
5. Bhandakkar, V.D.(2012), The study of molecular interaction in liquid mixtures using ultrasonic technique. *IOSR J.Appl.Phys.*,1(5), 38-43.
6. Bedare,G.R.,Bhandakkar, V.D. and Suryavanshi,B.M(2013), Acoustic behavior of cinnamaldehyde in polar and non-polar liquids at 298K., *Der Chemica sinica*, 4(1),97-101.
7. Mistry,A.A., Bhandakkar, V.D. and Chimankar,O.P.(2012), Acoustic studies on ternary mixture of toluene in cyclohexane at 308K using ultrasonic technique. *J.Chem.&Pharm.Res*, 4(1),170-174.
8. Bhandakkar, V.D. and Chimankar,O.P. and Mistry,A.A., *Pelagia Res.Lib.*,**2011**,2(6), 7076.